



Novel antigens used to detect cell-mediated immune responses over time in *Mycobacterium avium* subsp. *paratuberculosis* infected cattle

Mikkelsen, Heidi; Aagaard, Claus; Nielsen, Søren Saxmose ; Jungersen, Gregers

Publication date:
2009

[Link back to DTU Orbit](#)

Citation (APA):

Mikkelsen, H., Aagaard, C., Nielsen, S. S., & Jungersen, G. (2009). *Novel antigens used to detect cell-mediated immune responses over time in Mycobacterium avium subsp. paratuberculosis infected cattle*. Poster session presented at 8th Elsinore Meeting on infection immunity, Helsingør, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Novel antigens used to detect cell-mediated immune responses over time in *Mycobacterium avium* subsp. *paratuberculosis* infected cattle

Heidi Mikkelsen^{1,2}, Claus Aagaard³, Søren Saxmose Nielsen² and Gregers Jungersen¹

¹National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark; ²Faculty of Life Sciences, University of Copenhagen, Frederiksberg, Denmark, ³State Serum Institute; Copenhagen, Denmark
Contact: heimi@vet.dtu.dk

Background

Paratuberculosis is a chronic, granulomatous enteric infection caused by *Mycobacterium avium* subsp. *paratuberculosis* (MAP) in ruminants. Early-stage MAP infection can be detected using diagnostics for cell mediated immune responses, e.g. the whole blood interferon gamma (IFN- γ) test. Available IFN- γ tests are using purified protein derivatives of MAP (PPDj) which are crude products consisting of undefined antigens with possible cross reactions toward other environmental bacteria.

Objective

The objective was to optimize the IFN- γ test using different types of novel antigens for stimulation to determine if some antigens could be excluded or combined.


Methods

Fourteen novel antigens were selected for testing.


Blood samples were collected 3 times with 4 and 5 week intervals from the same 30 heifers 15-24 months of age in a herd with known MAP infection.

IFN- γ test protocol:

1) Stimulation of whole blood with MAP antigens




2) IFN- γ detection by ELISA



Conclusions

- PPDj stimulations tested approximately 50% of the animals as test positive at the two first samplings, and less than 30% on the third sampling.
- Ag85b tested 50% of the animals positive at all three samplings.
- The groups of latency proteins and secreted proteins tested approximately 40% of the animals test positive. Latency proteins showed the most consistent test results.
- ESAT-6 family proteins gave fluctuating test responses.

Results



IFN- γ responses = IFN- γ produced in response to specific antigen - IFN- γ produced in response to PBS-stimulation.

Interpretation

Cut-offs used to discriminate test-positives and test-negatives for each antigen was based on samples from 60 heifers 15-24 months of age from a non-infected herd.